
DSC 140A - Discussion 03

Problem 1.

Recall that the regularized least squares risk is

$$\tilde{R}(\vec{w}) = \frac{1}{n} \sum_{i=1}^n (\vec{w} \cdot \vec{\phi}(\vec{x}^{(i)}) - y_i)^2 + \lambda \|\vec{w}\|^2$$

Show that

$$\tilde{R}(\vec{w}) = \frac{1}{n} (\vec{w}^T \Phi^T \Phi \vec{w} - 2\vec{w}^T \Phi^T \vec{y} + \vec{y}^T \vec{y}) + \lambda \vec{w}^T \vec{w},$$

where Φ is the matrix whose i th row is $\vec{\phi}(\vec{x}^{(i)})$, and where $\vec{y} = (y_1, \dots, y_n)^T$.

Problem 2.

In lecture, we defined a kernel function to be a function k which computes the dot product of vectors after they are mapped to some high-dimensional space. The useful thing about kernel functions is that they allow us to compute these dot products without actually mapping vectors them to the high-dimensional space, which can be costly. In this problem, we will consider the the 2nd-order *polynomial kernel*, defined to be

$$k(\vec{x}, \vec{x}') = (1 + \vec{x} \cdot \vec{x}')^2.$$

Let $\vec{\phi}(\vec{x}) : \mathbb{R}^3 \rightarrow \mathbb{R}^{10}$ be the mapping:

$$\vec{\phi}(\vec{x}) = (1, x_1^2, x_2^2, x_3^2, \sqrt{2}x_1, \sqrt{2}x_2, \sqrt{2}x_3, \sqrt{2}x_1x_2, \sqrt{2}x_1x_3, \sqrt{2}x_2x_3)^T,$$

where x_1, x_2, x_3 are the components of the input vector, \vec{x} . That is, $\vec{\phi}$ is a feature map which maps a vector into a higher-dimensional space.

Show that $k(\vec{x}, \vec{y}) = \vec{\phi}(\vec{x}) \cdot \vec{\phi}(\vec{y})$. That is, that k indeed computes the inner product of vectors in the higher-dimensional space.